



A comparison of ground-based count methods for quantifying seed production in temperate broadleaved tree species

Clara Tattoni¹ · Francesco Chianucci² · Marco Ciolli³ · Carlotta Ferrara² · Luca Marchino² · Michele Zanni⁴ · Paolo Zatelli³ · Andrea Cutini²

Received: 8 July 2020 / Accepted: 16 November 2020 / Published online: 26 January 2021
© The Author(s) 2021

Abstract

• **Key message** Litter trap is considered the most effective method to quantify seed production, but it is expensive and time-consuming. Counting fallen seeds using a quadrat placed on the ground yields comparable estimates to the litter traps. Ground quadrat estimates derived from either visual counting in the field or image counting from quadrat photographs are comparable, with the latter being also robust in terms of user sensitivity.

• **Context** Accurate estimates of forest seed production are central for a wide range of ecological studies. As reference methods such as litter traps (LT) are cost- and time-consuming, there is a need of fast, reliable, and low-cost tools to quantify this variable in the field.

• **Aims** To test two indirect methods, which consist of counting the seeds fallen in quadrats.

• **Methods** The trial was performed in three broadleaved (beech, chestnut, and Turkey oak) tree species. Seeds are either manually counted in quadrats placed at the ground (GQ) or from images acquired in the same quadrats (IQ) and then compared against LT measurements.

• **Results** GQ and IQ provide fast and reliable estimates of seeds in both oak and chestnut. In particular, IQ is robust in terms of user sensitivity and potentially enables automation in the process of seed monitoring. A null-mast year in beech hindered validation of quadrats in beech.

• **Conclusion** Quadrat counting is a powerful tool to estimate forest seed production. We recommend using quadrats and LT to cross-calibrate the two methods in case of estimating seed biomass. Quadrats could then be used more routinely on account of their faster and simpler procedure to obtain measurements at more spatially extensive scales.

Keywords Mast seeding · Acorns · Nuts · Masting · Litter trap

1 Introduction

Mast seeding, also called masting, is the variable, intermittent production of large seed crops, which is a typical reproductive strategy of many wind-pollinated species. Masting events have

cascading effects on the overall ecosystem functioning. For instance, the associated resource pulses are relevant for the population dynamics of seed consumers like rodents (Elkinton et al. 1996; Zwolak et al. 2016), roe deer and wild boar (Bisi et al. 2016; Canu et al. 2015; Cutini et al. 2013; Jackson, 1980), brown bear (Ciucci et al. 2014; Tattoni et al. 2015), many bird species (Degange et al. 1989; Hannon et al. 1987; Szymkowiak and Kuczyński 2015; Czeszczewik and Walankiewicz 2016; Soler et al. 2017; Tattoni et al. 2019; Fležar et al. 2019; Szymkowiak and Thompson 2019), and insects (Bogdziewicz et al. 2018). Some forest tree seeds are also used as human or animal food supply (Tattoni et al. 2017); hence, their availability can provide an income in some rural area or add recreational value to forests (Riccioli et al. 2018). As masting determines seedling establishment and recruitment, it also plays a key role in forest management (Ascoli et al. 2015; Cutini et al. 2015; Chianucci

Handling Editor: Andreas Bolte

Contributions of the co-authors Conceptualization: FC; methodology: FC; formal analysis: CT; investigation: FC, MZ, CT, MC, PZ; writing—original draft: CT, FC; writing—review and editing: FC, MZ, CT, MC, PZ; supervision: AC; funding acquisition: AC.

✉ Clara Tattoni
clara.tattoni@gmail.com

Extended author information available on the last page of the article

et al. 2019a, b). Understanding masting is therefore crucial to improve the knowledge on population dynamics, assess present and future ecosystem resilience, and design adaptive forest management strategies (Cutini et al. 2013; Wagner et al. 2010).

Currently, there are some issues in studying mast seeding. Firstly, the definition of years with mast seeding (mast year) is controversial, and there are different methods to classify a mast year (Cutini et al. 2013; LaMontagne and Boutin 2009). Secondly, masting can be defined using different qualitative (e.g., Nussbaumer et al. 2018) or quantitative (e.g., Cutini et al. 2013; Koenig and Knops 2014) measurements and methods. While categorical data allow to harmonize information for wider-scale comparisons (Bajocco et al. 2020; Nussbaumer et al. 2018; Vacchiano et al. 2017), this can limit the understanding on the ‘real’ seed availability—i.e., the same species may exhibit a mast year, but with significantly different average seed production depending on the stand age, structure, and management (Cutini et al. 2015). Therefore, studies on quantitative data of seed production (e.g., number of seeds, seed biomass) can avoid the ambiguity of determining a mast year. However, the challenge in quantitatively measuring seed production is that available methods are time-consuming and expensive, which limits larger-scale application of quantitative approaches. As direct tree measurements are also impractical due to the height and density of forest canopies, ground-based methods have been frequently used to assess seed production (Perry et al. 1999).

Currently, two main ground methods have been used for quantifying seed production: litter traps and visual counting of seeds on the trees. The litter trap method (LT) has been used in the majorities of studies on forest seed production: examples of sampled species include beech (Bajocco et al. 2020; Chianucci et al. 2019a, b; Cutini et al. 2013), chestnut (Bisi et al. 2016; Cutini 2000, 2001), many oak species (Bogdziewicz et al. 2018; Chianucci et al. 2019a), but also conifers (Mencuccini et al. 1995). The LT is considered the most accurate method (Gea-Izquierdo et al. 2006; Perry et al. 1999), but it is limited by the cost and the time needed to either collect data (which requires frequent—generally biweekly—sampling) or process the litter in the laboratory (which requires separating litterfall by species and by main components—leaf, woody, reproductive parts). As an alternative to LT, indirect methods have also been tested, which are often based on visual counting of seeds while still on the trees. These surveys are undoubtedly faster and cheaper than LT, but they are limited by the subjectivity of the measurements, which are also not replicable, and the difficulty to apply them to tall trees, particularly those with small seed size or in situation of high tree density and canopy closure (Perry et al. 1999). In addition, visual surveys are also generally unable to yield a quantitative estimate of seed production.

Recently, a method based on counting the number of seeds after their falling on the ground was proposed by Touzot et al. (2018). The method can be considered a

floor-level variant of LT, but it has the further advantage of reducing the time and cost of field and laboratory working, being, therefore, more flexible, and allowing larger-scale field deployment compared with (fixed) litter traps. However, the method was tested only on oak forests, which are characterized by relatively large (and thus easily detectable) seed size on the ground. Therefore, more experiments are needed to sample tree seeds with different shape and size.

In this study, we tested two indirect ground-based methods to yield quantitative measurements (number) of seeds, which were compared against reference LT measurements. The first method was based on counting seeds on the ground in quadrats of known area, soon after seed fall (hereafter “ground quadrats”, abbreviated GQ); the method was similar to the ground plot method proposed by Touzot et al. (2018). The second was an image-based ground counting method (hereafter “image quadrats”, abbreviated IQ). Downward images of the forest floor were collected in the same quadrats used for GQ, soon after GQ counting, and then inspected to count the number of seeds. Image-based approaches for seed counting have been already proposed for agricultural crops (e.g., Mussadiq et al. 2015; Tańska et al. 2018). However, the existing solutions were able to retrieve seeds in an artificial homogeneous background. Unlike monoculture crops, the forest floor can contain litter layers, dead fallen leaves, coarse woody debris, bare ground rocks, which creates a complex background against which to detect seeds. In addition, the period of seed fall often coincides with leaf fall, and thus, fallen leaves can obscure seeds in the ground, hindering the image-based counting at the floor. Hence, the image analysis of tree seed at the ground must deal with the higher heterogeneity and complexity of the forest floor.

The trial was performed in three most diffuse broadleaves forest tree species in Italy. The ground quadrats estimates were first calibrated against benchmark values obtained by litter traps from a network of permanent plots (Chianucci et al. 2019a). Further ground quadrats measurements were then performed and compared with image quadrats, which were analyzed to test the reliability of the image-based counting of seeds. Our specific questions are.

1. Are quadrat seed counts comparable with LT?
2. Are IQ seed counts comparable with GQ?
3. Is IQ robust in terms of user sensitivity?

2 Material and methods

2.1 Study area

The study was performed in broadleaved forests sampled from six stations in central-Northern Italy (Fig. 1). Broadleaved forests in Italy represent around 76% of the national forest

surface (Tabacchi et al. 2007). We sampled pure forest stands of three most widespread broadleaved tree species in Italy: beech (*Fagus sylvatica* L.; 9.9% of the national forest surface; Tabacchi et al. 2007), Turkey oak (*Quercus cerris* L. 9.6%) and chestnut (*Castanea sativa* Mill. 7.5%). The distribution of the studied forests allowed to sample a very diverse range of environmental conditions, which is representative of the natural conditions of the sampled forest tree species.

2.2 Data collection

Field sampling was carried out in the stations during the period 16 October (day of year (doy) 289)–28 November 2019 (doy 332).

In two stations (AR03; AR04), concurrent measures of seed production were obtained with the three tested methods (LT,

GQ, IQ). Nine ground quadrats and litter trap measurements were taken for comparison between the three methods in pure forest stands where litter traps were installed, with quadrats placed close from litter traps; measurements were repeated 2–3 times in the sampled plots (Table 1). Quadrat counting and litterfall collection were performed on the same day for comparison between these methods (Tattoni and Chianucci 2020).

Additional quadrat measurements were collected in other pure stands of chestnut (AR01, 30 quadrats), beech (TN01, 52 quadrats), and oak (VT01 and AR02, 42 quadrats) by randomly placing quadrats in pure forest stand of the sampled species, with measurements repeated 1–2 times depending on the species (Table 1). Seeds were then removed from the traps/quadrats at the end of each sampling session. Table 1 lists the number of

Fig. 1 Study area. Location of the six stations where seed production of temperate broadleaved tree species was sampled with quadrats and seed traps. Green areas are forest coverage in Italy according to CORINE Land Cover level IV (CLC2006_CLC2000_V2018_20 seamless 100 m raster)




Table 1 Number of seed count measurements collected for pairwise comparison between ground seed counting methods

	Species	Pairs	Traps	Plots	Repetitions (day)
GQ, IQ, LT pairs	Chestnut	27	9	1	3 (289, 311, 330)
	Beech	36	9	2	2 (289, 330)
	Turkey oak	36	9	2	2 (289, 330)
Additional GQ, IQ pairs	Chestnut	30	-	1	1 (295)
	Beech	52	-	1	2 (308, 324)
	Turkey oak	42	-	1	2 (295, 332)

measurements collected for pairwise comparison between the seed counting methods.

2.3 Litter trap sampling

Litterfall was collected in nine 0.25-m² traps systematically distributed inside each permanent plot of AR03 and AR04 (Table 1). Litterfall was collected in the same days of quadrat collections in these plots, and then separated in the laboratory. The number of seeds per trap from LT was then

determined and used as a benchmark for comparison with GQ and IQ.

2.4 Ground quadrat sampling

A quadrat of 0.25 m² (the same size and shape of litter trap) was used to count the number of seeds in the ground (Fig. 2). In case of comparison with LT, the quadrat was placed north from the trap, at a distance within 1 m from it. The number of seeds was then manually counted in the field within each quadrat by a single observer (see Fig. 2).

Fig. 2 Examples of seed counting methods used to assess seed production in temperate broad-leaved tree species. A ground quadrat (0.25 m²) placed in beech **a**, Turkey oak **b**, chestnut **c** plots. A downward looking camera to collect quadrat images close to a litter trap **d**



2.5 Image-based quadrat sampling

We used a digital single lens reflex (Nikon D90) camera (Sendai Nikon Corp., Otawara, Tochigi, Japan) equipped with a Nikkor 18–200-mm lens locked to 18 mm to collect images of the ground quadrats, which were acquired during the same collection dates of LT (when available) and GQ. The camera was oriented perpendicular to the ground using a leveled tripod with a central column inclinable at 90° zenith angle (Vanguard Alta Pro 263AT, Vanguard World, China). An image centered on each ground sampled quadrat was then acquired (see Fig. 2), immediately after each GQ counting. Images were then downloaded, and the seeds manually counted on screen using ImageJ software Version 1.51 (Rasband 2018; Schneider et al. 2012). The estimated number of seeds from images was compared with those obtained from LT (when available) and GQ. Four images were disregarded because of low quality. A total of 219 images of the three species were analyzed (Table 1).

To assess the robustness of the IQ method in terms of its sensitivity to user subjectivity, three users further counted the number of seeds from three image sets, each consisting of 50 randomly selected images for each sampled species from the 219 images acquired. The user experience with the method ranged from beginners with no prior knowledge of seed counting to experts well trained with the IQ method. The results were compared in terms of the number of seeds, their standard deviations, and species-specific differences.

Table 2 summarizes the procedures of the three methods, including a comparison in terms of materials, costs, and number of operators needed. The estimation of costs should be interpreted with care, most of the laboratory equipment may be already available in most research centers, as well as the cameras, that can be used for multiple projects. Reflex cameras may be replaced by smartphone with good hardware in some cases, but this also needs testing. In Italy, the placement of an LT in the forest requires a formal authorization, but other countries may have different rules.

2.6 Statistical analyses

Firstly, we checked the difference in seed counts between the three methods using the Wilcoxon paired test (Zuur et al. 2010). We then compared seed counts using quadrats (GQ, IQ) against reference measurements using LT by fitting regression models. The influence of species on the estimated number of seeds by the counting methods was assessed using ANCOVA. Given that seed count data are usually non-normally distributed and often characterized by many zeros (Touzot et al. 2018; O'Hara and Kotze 2010), we adopted a zero inflated modeling approach. Several models were tested: Zero inflated Poisson (ZIP), Zero Inflated Negative Binomial (ZINB), Zero Altered Poisson (ZAP), Zero Altered Negative Binomial (ZANB) as named

by Zuur et al. (2009a). The last two models, known also as hurdle models, are similar to the zero-inflated ones, but more flexible about the zero modeling. We used odds ratio (OR) to interpret the model outputs because OR quantifies the strength of the association between two events and supports meaningful ecological interpretation (Keating and Cherry 2004; Rita and Komonen 2008). To consider the influence of seasonality on the fitting, we included the month of sampling as covariate of the regression. We then selected the best models using Akaike information criterion (AIC) and log likelihood.

We compared also quadrat counts between QG and IQ by fitting a linear regression to evaluate whether IQ are consistent with QG measurements. Finally, we compared the accuracy of IQ counts by different users, to evaluate the sensitivity of the method. ANOVA was used to assess the differences in seed counts by users independently for each species. The statistical analysis was performed with R and Rstudio interface (R Development Core Team 2011; R Studio 2015).

3 Results

Count data for all species and methods showed a non-normal, skewed distribution with many zeros and some outliers (especially for beech acorns), as illustrated in the box plots of Fig. 3a. The average number of seeds per trap measured from LT ranged between 0 and 42 in chestnut (average number \pm standard deviation 7.11 ± 11.01), between 0 and 8 in Turkey oak (2.23 ± 2.48), and between 0 and 7 in beech (1.00 ± 1.51).

The relatively high standard deviations were due to a high number of records without seeds: the proportion of zeros was 34.4% for IQ, 26.8% for GQ, and 44.4% for LT. The three species showed a different seasonality in seed falling (Fig. 3b): beech nuts fell mainly at the mid-end of November, while acorns and chestnuts fell mainly in October.

Wilcoxon paired test showed that seed counts in GQ and LT were not significantly different ($p > 0.05$), while the same test yielded a significant difference in medians between methods LT-IQ ($p < 0.05$) and GQ-IQ ($p < 0.05$). We also performed an ANCOVA test to the data, and we found a significant effect of species on calculated seed number from the different methods ($p < 0.05$).

3.1 Are ground quadrat and image quadrat seed counts comparable with litter trap counts?

3.1.1 GQ VS LT

To predict the number of seeds in the traps (LT) from ground quadrats (GQ) or image quadrats (IQ), we fitted different models using species and the month of collection as random effects.

Table 2 Comparison of the three methods in terms of equipment, procedures, costs, and operators needed. Costs are to be considered a tentative estimation; they may vary a lot according to the availability of cameras and laboratory equipments of research units. Travel expenses are calculated in case of the sampling site located within 100 km and a day in the field is about 6 working hours

	Litter traps (LT)	Ground quadrats (GQ)	Image quadrats (IQ)
Materials	Litter trap; bags; fan-forced oven; sieve, scale, and lab tools	Frame 50 × 50 cm	Reflex camera and lens; tripod; frame 50 × 50; SD card; laptop
Collection process	Installation of the LT; seed collection; separation of reproductive parts from total litter; drying of the seeds in oven (in case of biomass determination); seed count	Positioning of the frame on the ground; visual on field seed count	Positioning of the frame on the ground; tripod/camera setup; picture shooting; images download; visual on screen seed count
Storage	Physical storage of the seeds (not permanent)	–	Electronic storage of the images (permanent)
Permissions	Authorizations are required to place LT in the forest	No authorizations required	No authorizations required
Location	LT are installed in fixed locations	Frames are portable and can be easily moved	Frames and cameras are portable and can be easily moved
Operators	1–2	1	1
Estimated time per sample	2 h per trap (collection, separation, count) + 24 h in case of overdrying (seed biomass)	6 min per quadrat	6 min per quadrat (2 min shooting, up to 4 min for image download and on screen counting)
Equipment costs	250 Euros for each trap In case of biomass determination: 5000 Euro for laboratory equipments (fan-forced oven, digital scale, storage)	20 euro per frame	2000 euros for a reflex camera, lens, SD cards + 20 euros per frame
Costs per sample (Euros)	2	< 1	< 1
Samples per day (N)	18–25	40	40
Travel cost (Euros)	35	35	35




Fig. 3 **a** Boxplot showing the number of seeds collected in temperate broadleaved forest with the three methods: IQ (image quadrats), GQ (ground quadrats), and LT (litter traps). **b** Boxplot showing the number

of seeds per species and month of collection. The y-axis of both plots was cut at 50 to improve readability, 27 records of beech counts over 50 are not shown

The best model between GQ and LT was a zero altered negative binomial ZANB model, where the count model fitted a negative binomial distribution between the seed counts in quadrats and the seeds counted in the traps with interaction species (traps*species), while the zero model used species and month as covariates (more details about model selection are reported in Appendix Table 3). Odds ratio shows that in this model, the baseline of having a positive count vs zero is 1.19. The odds of detecting some seeds increase significantly in November and decrease for Turkey oak and beech (0.4 and 0.18, respectively). The positive count model explanation is that if there is any seed (positive counts), the average count is 2.27 seeds, but in case of beech and Turkey oak, the average count decreased, compared with the baseline chestnut, by 0.19 and 0.88, respectively (Appendix Table 5). The plot of predicted versus fitted curves (Fig. 4) of the count model shows the different fits of the model according to the species and month.

3.1.2 IQ VS LT

The same approach was used to find the best model predicting the relationship between IQ and LT. The best fitting model was a ZINB model negative binomial distribution between the seed counts in images and the seeds counted in the traps with interaction species, while the zero-model used species and month as covariates (more details of model selection and model coefficients are available in the Appendix section, Appendix Tables 3 and 4). The best fitting model for the count part was the same as the previous model (GQ VS LT); there was the same combination of covariates and similar coefficients for the count part. The zero part model had the same combination of covariates (species and month) but fitted a zero inflated negative binomial distribution instead of the zero altered negative binomial. The odds ratio for the count part was comparable with the above case. The baseline




Fig. 4 Observed and predicted values of the number seeds in litter traps (LT) according to the best ZANB model (formula: $LT \sim GQ * species_{species} + month$). **a** Count part: the estimate seed production in LT from seeds counted in ground quadrats (0.25 m^2) is different for

the three sampled species. **b** Zero model: the probability of having a positive seed count (instead of a zero) is higher in November (month 11) for all species and it is different across species




Fig. 5 Observed and predicted values of the number seeds in litter traps (LT) according to the best ZINB model (formula: $LT \sim IQ * species_{species} + month$). **a** Count part: the estimate seed production in LT from seeds counted in ground quadrats (0.25 m^2) is different for

the three species of temperate broadleaved trees. **b** Zero model: the probability of having a positive seed count (instead of a zero) is higher in November (month 11) for all species and it is different across species

odds of having a positive count vs zero is 1.19; it increased in November and decreased in the case of Turkey oak and beech (0.18 and 0.04, respectively). The positive count model has the chestnut as a baseline and the average count is 2.19 and decreased for beech and Turkey oak compared with chestnut by 0.19 and 0.88, respectively. See plots of Fig. 5 and Appendix Table 8.

3.2 Are image quadrats counts comparable with ground quadrats?

The whole dataset was used to assess the relationship between the seeds counted in the image (IQ) and the seeds counted in the fields (GQ). Due to the nature of these counts, we expected a linear correlation between IQ and GQ, so we tested several




Fig. 6 Observed values and regression line of seed counted in quadrats (0.25 m^2) predicted from seeds counted in images for three temperate broadleaved tree species (formula: $IQ = -0.57 + 0.86 \text{ GQ}$, adjusted R -squared: 0.96)

linear models with different covariates and interactions. The best model with lowest AIC was the simplest one, a linear regression between IQ and GQ (Fig. 6, R squared = 0.96, intercept = -0.5654 , slope = 0.8634); more details about the tests and model selection can be found in Appendix Table 9.

3.3 Is the image quadrat method robust in terms of user sensitivity?

Three different operators counted the seeds in the same 50 images per species picked at random, for a total of 150 pictures. The mean counts ranged between 12.5 and 13.6 in chestnut, between 5.5 and 5.8 in Turkey oak, between 13.1 and 15.7 in beech. ANOVA indicated that counting seeds on a picture is a reliable method, which is insensitive to users and their previous experience in counting seeds ($p = 0.9$).

4 Discussion

The main result of the study is that the quadrat counting, both IQ and GQ, can be used to quantify the number of seeds as compared with litter traps (LT). In fact, the two best models had an almost identical fitting for the count part, concerning the coefficients, their significance and overlapping predicted values (Figs. 4a and 5a). The models correctly predicted a higher number of chestnut seeds compared with Turkey oak and beech. Chestnut and oak coefficients were significant in both models, while they are not significant in beech. Despite fitting slightly different distributions (zero inflated for GQ and zero altered for IQ), the probability of zero counts is similar across models (Figs. 4b and 5b). The comparison also indicated a different performance of the method according to the sampled species. The results are in accordance with the findings of Touzot et al. (2018) for sessile oak (*Quercus petraea*) acorns. Unlike the previous study, in which four 0.25 m^2 quadrats were used for comparison with a 20-m^2 trap, we also demonstrated that using a single quadrat of the same size allowed a reliable estimate compared with a single litter trap; this allows reducing the sampling efforts per trap while simultaneously allowing increasing the number of quadrat samples across a site. This is an advantage per se of the GQ/IQ method, as the establishment of traps and monitoring of seed using LT is cost and time consuming, which limits their usage at the plot scale where they are permanently installed prior to authorization.

Sampling in November increased the probability of finding seeds for all the three species considered in this study. On top of that, late autumn is a time of the year when the herbal understorey vegetation is at its minimum facilitating the ground sampling with IQ and GQ. In the case of a thick litter of leaves, the operator can move the camera or remove some of the leaves to take a better image.

With reference to beech, results indicated a lower number of seeds and higher probability of not finding any, compared with the other species. We attributed such outcome to the combined effect of the relatively small size of beech nuts and the low fruiting production observed in 2019 (about three-fourth of the quadrats have zero seeds), which likely complicated the detection of (low number of small) seeds from quadrats. While Touzot et al. (2018) observed that no seeds in the quadrats correspond to very low fruiting levels, we speculate that the agreement between GQ and LT would likely increase in case of high fruiting (mast) years in beech when the production may likely reach more than 150 nuts/m^2 (Burschel et al. 1964) and up to over 300 nuts/m^2 (Schmidt 2006).

An additional improvement of the GQ method in this species would be collecting the litter in the quadrat and then separating it in the laboratory, to reduce the probability of missing counts of small seeds in the field. This procedure is likely to help counting small-sized seeds of other trees, such as ashes and hornbeams (Czeszczewik et al. 2020).

Zero inflated model approach was necessary to account for the high number of zeros in the data set, which is typical of seed fall, particularly in low fruiting years (Calama et al. 2011; Touzot et al. 2018). We advise the need for future investigation, in a multiannual time-scale, to evaluate the model fitting of seed counts in higher fruiting (particularly mast) years.

With reference to quadrat counting, we demonstrated that IQ is statistically comparable with GQ. The outcome further extends the applicability of quadrat sampling, since collecting images is faster than ground quadrat counting, allowing a higher number of samples to be performed. In addition, IQ has further advantages compared with GQ: firstly, images are permanent records, which could be inspected to check for data quality (Chianucci et al. 2019b). Secondly, IQ is robust and provides similar estimates of seed numbers, irrespective of the users and their previous knowledge on seed counting. Thirdly, images can also be re-analyzed to test different methods to count seeds from images. In this line, some image classification methods could also be developed and compared against the manual counting from seed images. At the moment, the available tools in image classification could offer promising solution such as object-based image analysis (OBIA) or with convolutional neural network (CNN). However, these methods have been tested only in laboratory conditions and for agricultural crops, not for forest seeds in a natural, spectrally complex background. Both solutions require some training of the algorithms for classification (OBIA) or for detection and recognition of a search image of the seeds (CNN).

A potential limitation of IQ is that seed fall typically occurs simultaneously with leaf fall, and therefore, some fallen leaves in the ground could cause underestimation of seeds captured by images, as compared with LT and GQ. Although the observed differences between IQ and GQ were not statistically significant,

this issue further supports the need to set a proper timing for quadrat counting, possibly increasing the frequency of sampling.

To sum up, quadrat counting represents a simple, robust, cheap and flexible method to estimate tree seed number in the field. From a practical viewpoint, considering the long-term perspective of seed (and masting) dynamics, we suggest to calibrate quadrats counting with measurements obtained from LT in calibration plots (when available); this would also allow relating seed number with either continuous (seed mass) or categorical (low, medium, high fruiting, or other scales) measurements of seed in reference plots. Both GQ and IQ could be then used more routinely for indirect measurements and monitoring of seed production on larger areas.

As we focused on pure methodological differences, we did not apply a probabilistic sampling scheme as it would be recommended for, e.g., estimating seed production at the tree, stand, or site level. Therefore, future investigations are required to assess the sampling efforts (scheme and number of samples) required to obtain statistically representative estimates of seed at different sampling domains. For the aim of this study, we sampled pure stands, but we believe that the method can be applied also in mixed forests, as the seeds of each species are recognizable with a high degree of accuracy by a trained operator.

5 Conclusions

Field estimates of seed production are essential for a wide range of ecological studies. We demonstrated that ground quadrats are fast, simple, and reliable tools to assess seed production in broadleaf forest stands, being comparable with the more consuming litter trap method. Among the quadrats method, image quadrats hold great potential for future researches and applications, including the development of seed monitoring systems based on continuous camera, as already existing in other monitoring applications such as tree phenology (Chianucci 2020; Richardson et al. 2018). In this line, downward looking cameras could be installed to capture images at the forest floor for counting the number of seeds. This option would further reduce (or avoid) the efforts required for field surveys, simultaneously allowing an assessment of seed fall patterns at a finer, daily temporal scale.

Appendix

This section reports some supplemental information about the model selection.

Table organization and content:

Tables 3, 4, 5: GQ VS LT model selection, coefficients, odds ratio.

Table 3 Model selection for the relationship between seed production estimated with ground quadrats (GQ) and the reference litter trap method (LT) for the sampled tree species. AIC and log likelihood were used to select the best models (in italics)

Model	Vars. count model	Vars. zero model	AIC	LogLik	Pr (>Chisq)
ZIP	LT	LT	456.3	- 224.1	
ZIP	LT + species	LT + species	442.2	- 213.1	22.0
ZIP	LT	Species	445.3	- 215.6	5.0
ZIP	LT* species	Species + month	441.6	- 214.8	1.7
ZINB	LT	LT	387.5	- 188.8	52.0
ZINB	LT + species	LT + Species	386.6	- 187.3	2.9
ZINB	LT	Species + month	384.2	- 184.1	6.3
ZINB	LT* species	Species + month	368.8	- 173.4	21.4
ZAP	LT	LT	456.3	- 224.1	79.3
ZAP	LT + species	LT + species	447.6	- 217.8	12.7
ZAP	LT	Species + month	436.8	- 210.4	14.7
ZAP	LT* species	Species + month	440.7	- 214.4	7.9
ZANB	LT	LT	387.6	- 188.8	13.1
ZANB	LT + species	LT + species	382.5	- 182.2	64.2
ZANB	LT	Species + month	367.6	- 172.8	31.9
ZANB	<i>LT* species</i>	<i>Species + month</i>	364.6	- 171.3	9.0

Table 4 GQ VS LT model coefficients

Count model coefficients (truncated negbin with log link)				
	Estimate	Std. error	z	value Pr(> z)
Intercept	0.865	0.409	2.117	0.034 *
LT	0.052	0.023	2.255	0.024 *
T. oak	- 1.636	0.578	- 2.831	0.005 **
Beech	- 0.124	0.628	- 0.198	0.843
LT: T. oak	0.224	0.122	1.836	0.066
LT: beech	0.031	0.294	0.106	0.916
Log(theta)	- 0.259	0.610	- 0.425	0.671
Zero altered model coefficients (binomial with logit link)				
Intercept	0.174	0.601	0.290	0.772
T. oak	- 1.693	0.780	- 2.172	0.030 *
Beech	- 3.025	0.817	- 3.703	0.0002 ***
November	2.781	0.689	4.035	0.00005 ***

Table 5 Odds ratios (OR) of ZANB model GQ VS LT

Count model	OR	Zero model	OR
Intercept	2.375	Intercept	1.190
LT	1.054	T. Oak	0.184
T. Oak	0.195	Beech	0.049
Beech	0.883	November	16.131
LT: T. Oak	1.251		
LT: beech	1.032		

Tables 6, 7, 8: IQ VS LT model selection, coefficients, odds ratio.

Table 6 Model selection for the relationship between seed production estimated with image quadrats (GQ) and the reference litter trap method (LT) for the sampled tree species. Month refers to the month of sampling. AIC and log likelihood were used to select the best models (in italics)

Model	Vars. count model	Vars. zero model	AIC	LogLik	Pr(>Chisq)
ZIP	LT + species	LT + species	355.96	- 169.9814	2.72E-01
ZIP	LT	Species + month	361.91	- 174.9570	9.04E + 02
ZIP	LT* species	Species + month	350.49	- 165.2495	5.13E + 01
ZINB	LT	LT	324.84	- 157.4203	8.90E + 02
ZINB	LT + species	LT + species	308.01	- 145.0094	4.63E + 00
ZINB	LT	Species + month	302.31	- 144.1550	2.55E + 04
<i>ZINB</i>	<i>LT* species</i>	<i>Species + month</i>	<i>302.17</i>	<i>- 140.0867</i>	<i>6.69E + 03</i>
ZAP	LT	LT	377.45	- 184.7261	7.40E-11
ZAP	LT + species	LT + species	358.13	- 171.0651	7.11E + 00
ZAP	LT	Species + month	363.47	- 175.7358	3.66E + 02
ZAP	LT* species	Species + month	355.88	- 167.9446	6.34E + 02
ZANB	LT	LT	323.90	- 156.9507	2.64E + 01
ZANB	LT + species	LT + species	312.64	- 147.3223	9.97E + 01
ZANB	LT	Species + month	309.92	- 147.9603	2.83E + 04
ZANB	LT* species	Species + month	311.83	- 144.9187	9.30E + 04

Table 7 IQ VS LT model coefficients and significance

Count model coefficients (negbin with log link)				
	Estimate	Std. error	z value	Pr(> z)
Intercept	0.785	0.246	3.187	0.001 **
LT	0.028	0.017	1.672	0.094
T. oak	- 1.339	0.607	- 2.207	0.027 *
Beech	0.571	0.542	1.054	0.292
LT: T. oak	0.198	0.110	1.792	0.073
LT: beech	- 0.018	0.187	- 0.097	0.922
Log(theta)	0.297	0.371	0.8	0.423
Zero-inflation model coefficients (binomial with logit link):				
Intercept	- 12.17	190.61	- 0.064	0.949
T. oak	21.74	198.07	0.110	0.913
Beech	23.40	198.07	0.180	0.906
November	10.93	53.85	- 0.203	0.839

Table 8 Odds ratios of model IQ VS LT

Count model	OR	Zero model	OR
Intercept	2.193	Intercept	5.167e-06
LT	1.029	T. oak	2.752e + 09
T. oak	0.262	Beech	1.460e + 10
Beech	1.771	November	1.795e-05
LT: T. oak	1.219		
LT: beech	0.982		

Tables 9 and 10: GQ VS IQ model selection and coefficients.

Table 9 Model selection for the relationship between seed production estimated with image quadrats IQ and the reference ground quadrats (GQ) method. AIC and log likelihood were used to select the best model (in italics)

Model	Variables	R-squared	df	AIC	LogLik	Pr(>Chisq)
<i>LM</i>	<i>GQ</i>	0.958	3	1121.83	- 557.91	
LM	GQ + species	0.957	5	1125.28	- 557.64	0.762
LM	GQ* species	0.958	7	1125.77	- 555.88	0.172
LM	GQ + month	0.958	4	1123.23	- 557.61	0.325
LM	GQ* month	0.958	5	1124.84	- 557.42	0.535
LM	GQ + month + species	0.958	6	1132.86	- 556.91	0.312

Table 10 Coefficients of linear model IQ VS GQ, adjusted R-squared: 0.9582

	Estimate	Std. error	t value	Pr(> t)
Intercept	- 0.5654	0.273	- 2.072	0.0395 *
GQ	0.863	0.012	68.909	2e-16 ***

List of Abbreviation

GQ Ground quadrats;

IQ Image quadrats;

LT Litter traps;

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Model abbreviations are reported according to Zuur et al (2019b). Formulas are reported according to R standards, where + means linear combination of variables and * means interaction. Month refers to the month of sampling

Acknowledgments Field assistance from the personnel of CREA, particularly Maurizio Piovosi, Leonardo Tonveronachi, Tessa Giannini, and Giovanbattista De Dato is greatly acknowledged. We dedicated this study to the memory of Alfredo Bresciani.

Funding Open access funding provided by Università degli Studi di Firenze within the CRUI-CARE Agreement. The study was financially supported by the research project “Monitoraggio della produzione di seme di specie forestali, rinnovazione naturale e relazioni con la fauna selvatica (Pasciona)” funded by the Foreste Casentinesi, Monte Falterona e Campigna National Park.

Data availability The dataset generated during the current study is available on Mendeley repository: <https://data.mendeley.com/datasets/cc6m499bmm>.

Compliance with ethical standard

Conflict of interest The authors declare that they have no conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing,

adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.


References

- Ascoli D, Vacchiano G, Maringer J, Bovio G, Conedera M (2015) The synchronicity of masting and intermediate severity fire effects favors beech recruitment. For Ecol Manage 353:126–135. <https://doi.org/10.1016/j.foreco.2015.05.031>
- Bajocco S, Ferrara C, Bascietto M, Alivernini A, Chirichella R, Cutini A, Chianucci F (2020) Characterizing the climatic niche of mast seeding in beech: Evidences of trade-offs between vegetation growth and seed production. Ecological Indicators:107139
- Bisi F, von Hardenberg J, Bertolino S, Wauters LA, Imperio S, Preatoni DG, Provenzale A, Mazzamuto MV, Martinoli A (2016) Current and future conifer seed production in the Alps: testing weather factors as cues behind masting. Eur J For Res 135:743–754. <https://doi.org/10.1007/s10342-016-0969-4>
- Bogdziewicz M, Steele MA, Marino S, Crone EE (2018) Correlated seed failure as an environmental veto to synchronize reproduction of masting plants. New Phytol 219:98–108. <https://doi.org/10.1111/nph.15108>
- Burschel P, Huss J, Kalbhenn R (1964) Natural regeneration of beech. Schriftenr. Forstl. Fak. Univ. Göttingen 34: 186 p [German]
- Calama R, Mutke S, Tomé J, Gordo J, Montero G, Tomé M (2011) Modelling spatial and temporal variability in a zero-inflated variable: the case of stone pine (*Pinus pinea* L.) cone production. Ecol Model 222(3): 606–618
- Canu A, Scandura M, Merli E, Chirichella R, Bottero E, Chianucci F, Cutini A, Apollonio M (2015) Reproductive phenology and conception synchrony in a natural wild boar population. Hystrix 26: <https://doi.org/10.4404/hystrix-26.2-11324>
- Chianucci F (2020) An overview of in situ digital canopy photography in forestry. Can J For Res 50(3):227–242
- Chianucci F, Ferrara C, Bertini G, Fabbio G, Tattoni C, Rocchini D, Corona P, Cutini A (2019a) Multi-temporal dataset of stand and canopy structural data in temperate and Mediterranean coppice forests. Ann For Sci 76:80. <https://doi.org/10.1007/s13595-019-0862-7>

- Chianucci F, Ferrara C, Pollastrini M, Corona P (2019b) Development of digital photographic approaches to assess leaf traits in broadleaf tree species. *Ecol Ind* 106:105547
- Ciucci P, Tosoni E, Di Domenico G, Quattrociochi F, Boitani L (2014) Seasonal and annual variation in the food habits of Apennine brown bears, central Italy. *J Mammal* 95:572–586. <https://doi.org/10.1644/13-MAMM-A-218>
- Cutini A (2000) Biomass, litterfall and productivity in chestnut coppices of various age at Monte Amiata (Central Italy). *Ecol Mediterr* 26:33–41
- Cutini A (2001) New management options in chestnut coppices: an evaluation on ecological bases. *For Ecol Manage* 141:165–174
- Cutini A, Chianucci F, Chirichella R, Donaggio E, Mattioli L, Apollonio M (2013) Mast seeding in deciduous forests of the northern Apennines (Italy) and its influence on wild boar population dynamics. *Ann For Sci* 70:493–502. <https://doi.org/10.1007/s13595-013-0282-z>
- Cutini A, Chianucci F, Giannini T, Manetti M, Salvati L (2015) Is anticipated seed cutting an effective option to accelerate transition to high forest in European beech (*Fagus sylvatica* L.) coppice stands? *Ann For Sci* 72:631–640. <https://doi.org/10.1007/s13595-015-0476-7>
- Czeszczewik D, Walankiewicz W (2016) Ecology and biology of birds in the Białowieża Forest: a 40-year perspective. *For Res Pap* 77(4):332–340
- Czeszczewik D, Czortek P, Jaroszewicz ZK, Rowiński P, Walankiewicz W (2020) Climate change has cascading effects on tree masting and the breeding performance of a forest songbird in a primeval forest. *Sci Total Environ* 47:142084. <https://doi.org/10.1016/j.scitotenv.2020.142084>
- Degange AR, Fitzpatrick JW, Layne JN, Woolfenden GE (1989) Acorn harvesting by Florida scrub jays. *Ecology* 70:348–356. <https://doi.org/10.2307/1937539>
- Elkinton JS, Healy WM, Buonaccorsi JP, Boettner GH, Hazzard AM, Smith HR, Liebhold AM (1996) Interactions among gypsy moths, white-footed mice, and acorns. *Ecology* 77:2332–2342. <https://doi.org/10.2307/2265735>
- Fležar U, Costa C, Bordjan D, Jerina K, Krofel M (2019) Free food for everyone: artificial feeding of brown bears provides food for many non-target species. *Eur J Wildl Res* 65:1. <https://doi.org/10.1007/s10344-018-1237-3>
- Gea-Izquierdo G, Cañellas I, Montero G (2006) Acorn production in Spanish holm oak woodlands. *Investig Agrar Sist y Recur For* 15:339. <https://doi.org/10.5424/srfi/2006153-00976>
- Hannon SJ, Mumme RL, Koenig WD, Spon S, Pitelka FA (1987) Poor acorn crop, dominance, and decline in numbers of acorn woodpeckers. *J Anim Ecol* 56:197. <https://doi.org/10.2307/4809>
- Jackson J (1980) The annual diet of the roe deer (*Capreolus capreolus*) in the New Forest, Hampshire, as determined by rumen content analysis. *J Zool* 192:71–83. <https://doi.org/10.1111/j.1469-7998.1980.tb04220.x>
- Keating KA, Cherry S (2004) Use and interpretation of logistic regression in habitat-selection studies. *J Wildl Manage* 68:774–789. [https://doi.org/10.2193/0022-541x\(2004\)068\[0774:uaiolr\]2.0.co;2](https://doi.org/10.2193/0022-541x(2004)068[0774:uaiolr]2.0.co;2)
- Koenig WD, Knops JMH (2014) Environmental correlates of acorn production by four species of Minnesota oaks. *Popul Ecol* 56:63–71. <https://doi.org/10.1007/s10144-013-0408-z>
- LaMontagne JM, Boutin S (2009) Quantitative methods for defining mast-seeding years across species and studies. *J Veg Sci* 20:745–753. <https://doi.org/10.1111/j.1654-1103.2009.01068.x>
- Mencuccini M, Piussi P, Zanzi Sulli A (1995) Thirty years of seed production in a subalpine Norway spruce forest: patterns of temporal and spatial variation. *For Ecol Manage* 76:109–125. [https://doi.org/10.1016/0378-1127\(95\)03555-O](https://doi.org/10.1016/0378-1127(95)03555-O)
- Mussadiq Z, Laszlo HL, Gyuricza C (2015) Evaluation and comparison of open source program solutions for automatic seed counting on digital images. *Comput Electron Agric* 117:194–199. <https://doi.org/10.1016/j.compag.2015.08.010>
- Nussbaumer A, Waldner P, Apuhtin V, Aytar F, Benham S, Bussotti F, Eichhorn J, Eickenscheidt N, Fabianek P, Falkenried L, Leca S, Lindgren M, Manzano Serrano MJ, Neagu S, Nevalainen S, Pajtik J, Potočić N, Rautio P, Sioen G, Stakénas V, Tasdemir C, Thomsen IM, Timmermann V, Ukonmaanaho L, Verstraeten A, Wulff S, Gessler A (2018) Impact of weather cues and resource dynamics on mast occurrence in the main forest tree species in Europe. *For Ecol Manage* 429:336–350. <https://doi.org/10.1016/j.foreco.2018.07.011>
- O'Hara R, Kotze DJ (2010) Do not log-transform count data. *Methods Ecol Evol* 1:118–122. <https://doi.org/10.1111/j.2041-210x.2010.00021.x>
- Perry RW, Thill RE, Peitz DG, Tappe PA (1999) Effects of different silvicultural systems on initial soft mast production. *Wildl Soc Bull* 27:915–923. <https://doi.org/10.2307/3783646>
- R Development Core Team, R. R (2011) R: a language and environment for statistical computing. *Stat. Comput., R Foundation for Statistical Computing, R Found.* <https://doi.org/10.1007/978-3-540-74686-7>
- R Studio, (2015) RStudio: Integrated development environment for R
- Rasband WS (2018) ImageJ [WWW Document]. Natl. Institutes Heal. Bethesda, Maryland, USA. URL <https://imagej.nih.gov/ij/>, 1997–2018. Accessed 3.17.20
- Riccioli F, Marone E, Boncinelli F, Tattoni C, Rocchini D, Fratini, R (2018) The recreational value of forests under different management systems. *New For.* 1–16. <https://doi.org/10.1007/s11056-018-9663-3>
- Richardson AD, Hufkens K, Milliman T, Aubrecht DM, Chen M, Gray JM, Johnston MR, Keenan TF, Klosterman ST, Kosmala M, Melaas EK, Friedl MA, Froking S (2018) Tracking vegetation phenology across diverse North American biomes using PhenoCam imagery. *Sci Data* 5:1–24. <https://doi.org/10.1038/sdata.2018.28>
- Rita H, Komonen A (2008) Odds ratio: an ecologically sound tool to compare proportions. *Annales Zoologici Fennici* 45:66–72. <https://doi.org/10.5735/086.045.0106>
- Schmidt, W (2006) Temporal variation in beech masting (*Fagus sylvatica* L.) in a limestone beech forest (1981–2004). *Allg. Forst- u. J.-Ztg*: 177: 9–19 [German]
- Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image analysis. *Nat Methods* 9:671–675. <https://doi.org/10.1038/nmeth.2089>
- Soler R, Espelta JM, Lencinas MV, Peri PL, Martínez Pastur G (2017) Masting has different effects on seed predation by insects and birds in Antarctic beech forests with no influence of forest management. *For Ecol Manage* 400:173–180. <https://doi.org/10.1016/j.foreco.2017.06.014>
- Szymkowiak J, Kuczyński L (2015) Avoiding predators in a fluctuating environment: responses of the wood warbler to pulsed resources. *Behav Ecol* 26(2):601–608
- Szymkowiak J, Thomson RL (2019) Nest predator avoidance during habitat selection of a songbird varies with mast peaks and troughs. *Behav Ecol Sociobiol* 73(7):91
- Tabacchi G, De Natale F, Di Cosmo L, Floris A, Gagliano C, Gasparini P, Genchi L, Scrinzi G, Tosi V (2007) Forest cover estimate 2005 - Inventario Nazionale delle Foreste e dei Serbatoi Forestali di Carbonio. [WWW Document]. MiPAF - Corpo For. dello Stato - Isp. Gen. CRA - ISAFa, Trento. URL https://www.sian.it/inventarioforestale/jsp/home_en.jsp Accessed 3.4.20
- Tańska M, Buczek J, Jarecki W, Wasilewska A, Konopka I (2018) Grain morphology, texture and colour-related compounds of bread wheat cultivars in relation to cultivation regimes and growing location. *Zemdirbyste* 105:105–112. <https://doi.org/10.13080/z-a.2018.105.014>
- Tattoni C, Bragalanti N, Groff C, Rovero F (2015) Patterns in the use of rub trees by the Eurasian Brown Bear. *Hystrix Ital J Mammal* 26. <https://doi.org/10.4404/hystrix-26.2-11414>
- Tattoni C, Ianni E, Geneletti D, Zatelli P, Ciolli M (2017) Landscape changes, traditional ecological knowledge and future scenarios in the

- Alps: a holistic ecological approach. *Sci Total Environ* 579:27–36. <https://doi.org/10.1016/j.scitotenv.2016.11.075>
- Tattoni C, Chianucci, F (2020) Dataset of seed production in temperate broadleaved tree species using different sampling methods. *Mendeley Repo* . V2. <https://data.mendeley.com/datasets/cc6m499bmm/>. Accessed 27/10/2020
- Tattoni C, Soardi E, Prosser F, Zatelli P, Odasso M, Ciolli M (2019) Fruit availability for migratory birds: a GIS approach. *PeerJ* 7:e6394. <https://doi.org/10.7717/peerj.6394>
- Touzot L, Bel-Venner MC, Gamelon M, Focardi S, Boulanger V, Débias F, Delzon S, Saïd S, Schermer E, Baubet E, Gaillard JM, Venner S (2018) The ground plot counting method: a valid and reliable assessment tool for quantifying seed production in temperate oak forests? *For Ecol Manage* 430:143–149. <https://doi.org/10.1016/j.foreco.2018.07.061>
- Vacchiano G, Hackett-Pain A, Turco M, Motta P, Maringer J, Conedera M, Drobyshev I, Ascoli D (2017) Spatial patterns and broad-scale weather cues of beech mast seeding in Europe. *New Phytol* 215:595–608. <https://doi.org/10.1111/nph.14600>
- Wagner F, Rutishauser E, Blanc L, Herault C (2010) Effects of plot size and census interval on descriptors of forest structure and dynamics. *Biotropica* 42:664–671. <https://doi.org/10.1111/j.1744-7429.2010.00644.x>
- Zuur AF, Ieno EN, Elphick CS (2010) A protocol for data exploration to avoid common statistical problems. *Methods Ecol Evol* 1:3–14. <https://doi.org/10.1111/j.2041-210X.2009.00001.x>
- Zuur AF, Ieno EN, Walker NJ, Saveliev AA, Smith GM (2009a) Zero-truncated and zero-inflated models for count data. 261–293. https://doi.org/10.1007/978-0-387-87458-6_11
- Zuur AF, Ieno EN, Walker NJ, Saveliev AA, Smith GM (2009b) Mixed effects modelling for nested data. 101–142. https://doi.org/10.1007/978-0-387-87458-6_5
- Zwolak R, Bogdziewicz M, Wróbel A, Crone EE (2016) Advantages of masting in European beech: timing of granivore satiation and benefits of seed caching support the predator dispersal hypothesis. *Oecologia* 180:749–758. <https://doi.org/10.1007/s00442-015-3511-3>

Authors and Affiliations

Clara Tattoni¹  · Francesco Chianucci² · Marco Ciolli³ · Carlotta Ferrara² · Luca Marchino² · Michele Zanni⁴ · Paolo Zatelli³ · Andrea Cutini²

Francesco Chianucci
francesco.chianucci@crea.gov.it

Marco Ciolli
marco.ciolli@unitn.it

Carlotta Ferrara
carlotta.ferrara@crea.gov.it

Luca Marchino
luca.marchino@crea.gov.it

Michele Zanni
zanni.michele2@gmail.com

Paolo Zatelli
paolo.zatelli@unitn.it

Andrea Cutini
andrea.cutini@crea.gov.it

- ¹ DAGRI Dipartimento Di Scienze E Tecnologie Agrarie, Alimentari Ambientali E Forestali, University of Florence, Firenze, Italy
- ² CREA - Research Centre for Forestry and Wood, Arezzo, Italy
- ³ DICAM - Dipartimento Ingegneria Civile Ambientale, e Meccanica University of Trento, Trento, Italy
- ⁴ Department of Veterinary Medicine, University of Sassari, Sassari, Italy